

What do I need to know after each PLUVICTO dose?

1. Help reduce radiation exposure to yourself and others

Your body, blood, sweat, and urine give off radiation for a while after getting PLUVICTO. So, it is important to follow these instructions. For all these instructions, count **Day 1** as the day after you get PLUVICTO.



Stay hydrated

For 1 day drink a lot and urinate (pee) as much as you can



Stay more than 3 feet away from others

For 2 days from other adults

For 7 days from children under 18 or pregnant women



Sleep alone in a separate bedroom

For 3 days away from other adults

For 7 days away from children

For 15 days away from pregnant women



No sex at first, and then use effective birth control

For 7 days no sex

During your course of treatment and for 14 weeks after your last dose use effective birth control to avoid pregnancy



Use the bathroom carefully

Use a separate bathroom if possible, and wipe it down after you use it

For 2 days always sit while using the toilet
Use and flush toilet paper every time

For 3 days whenever anyone helps you in the bathroom, they should wear disposable gloves



Shower daily

For 7 days take at least one shower a day



Do separate laundry

During your course of treatment wash your laundry separately from others

Be aware that treatment may cause infertility

2. Be on the lookout for side effects during and after your treatment

Tell your doctor about any side effects you have because something may be done to help.

The most common side effects of PLUVICTO are:

- Tiredness
- Dry mouth
- Nausea (belly pain)
- Low red blood cell count (feeling weak or short of breath)
- Loss of appetite
- Changes in bowel movements (constipation or diarrhea)
- Vomiting (throwing up)
- Low blood platelet count (bleeding or bruising more easily than normal)
- Urinary tract infection
- Weight loss
- Abdominal pain (belly pain)

3. Plan your next bloodwork appointment with your doctor

You will need to have your bloodwork done again 1-2 weeks before the next time you get PLUVICTO.

For more information, please visit us at [PLUVICTO.com](https://www.pluvicto.com) or contact your doctor

Please see Important Safety Information on back.

 **PLUVICTO**[®]
lutetium Lu 177 vipivotide tetraxetan
INJECTION FOR INTRAVENOUS USE

Important Safety Information

What is PLUVICTO® (lutetium Lu 177 vipivotide tetraxetan)?

PLUVICTO is a radiopharmaceutical used to treat adults with an advanced cancer called prostate-specific membrane antigen-positive metastatic castration-resistant prostate cancer (PSMA-positive mCRPC) that:

- has spread to other parts of the body (metastatic), and
- has already been treated with other anticancer treatments

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about PLUVICTO?

Use of PLUVICTO involves exposure to radioactivity. Long-term, accruing radiation exposure is associated with an increased risk for cancer.

To minimize radiation exposure to others following administration of PLUVICTO, limit close contact (less than 3 feet) with household contacts for 2 days or with children and pregnant women for 7 days, refrain from sexual activity for 7 days, and sleep in a separate bedroom from household contacts for 3 days, from children for 7 days, or from pregnant women for 15 days.

PLUVICTO may cause serious side effects, including:

- **Low level of blood cell counts.** Tell your doctor right away if you develop any new or worsening symptoms, including:
 - Tiredness or weakness
 - Pale skin
 - Shortness of breath
 - Bleeding or bruising more easily than normal or difficulty stopping bleeding
 - Frequent infections with signs such as fever, chills, sore throat, or mouth ulcers
- **Kidney problems.** Tell your doctor right away if you develop any new or worsening symptoms, including passing urine less often or passing much smaller amounts of urine than usual

Before you receive PLUVICTO, tell your doctor if any of these apply to you:

- You have low level of blood cell counts (hemoglobin, white blood cell count, absolute neutrophil count, platelet count)
- You have or have had tiredness, weakness, pale skin, shortness of breath, bleeding or bruising more easily

than normal or difficulty stopping bleeding, or frequent infections with signs such as fever, chills, sore throat, or mouth ulcers (possible signs of myelosuppression)

- You have or have had kidney problems
- You have or have had any other type of cancer or treatment for cancer, as PLUVICTO contributes to your long-term cumulative radiation exposure
- You are sexually active as:
 - All radiopharmaceuticals, including PLUVICTO, have the potential to cause harm to an unborn baby
 - You should use effective contraception for intercourse during treatment with PLUVICTO and for 14 weeks after your last dose
 - PLUVICTO may cause temporary or permanent infertility

Before administration of PLUVICTO, you should drink plenty of water in order to urinate as often as possible during the first hours after administration.

The most common side effects of PLUVICTO include:

- Tiredness
- Dry mouth
- Nausea
- Low red blood cell count
- Loss of appetite
- Changes in bowel movements (constipation or diarrhea)
- Vomiting
- Low blood platelet count
- Urinary tract infection
- Weight loss
- Abdominal pain

These are not all of the possible side effects of PLUVICTO. Call your doctor for advice about side effects. You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

Please see full Prescribing Information.

